

PrimeGene Technical Data Sheet

Catalog Number:	101-17
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 31.0 kDa, a disulfide-linked homodimer of two 132 amino acid polypeptide chains.
Quantity:	5µg/25µg/1000µg
AA Sequence:	GITIPRNPGC PNEEDKNFPR TVMVNLSIHN RNTNTNPKRS SDYYNRSTSP WNLHRNEDPE RYPVSIWEAK CRHLGCINAD GNVDYHMNSV PIQQEILVLR REPPHCPNSF RLEKILVSVG CTCVTPIVHH VA
Purity:	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The ED ₅₀ as determined by inducing IL-6 secretion of murine NIH/3T3 cells is less than 7.5 ng/ml, corresponding to a specific activity of > 1.3 × 10 ⁵ IU/mg.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Endotoxin:	Less than 1 EU/µg of rHuIL-17 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in 4 mM HCl to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Human Interleukin-17

Human Interleukin-17A (IL-17A) is encoded by the IL17A gene located on the chromosome 6 and belongs to the IL-17 family that contains IL-17A, IL-17B, IL-17C, IL-17D, IL-17E and IL-17F. They have a similar protein structure, with four highly conserved cysteine residues critical to their 3-dimensional shape, but no sequence similarity to any other known cytokines. Interleukin 17 is a T cell-expressed pleiotropic cytokine that exhibits a high degree of homology to a protein encoded by the ORF13 gene of herpesvirus Saimiri. Mature IL-17 containing one potential N-linked glycosylation site. Both recombinant and natural IL-17 have been shown to exist as disulfide linked homodimers. At the amino acid level, IL-17 exhibits 63 % amino acid identity with mouse IL-17. High levels of human IL-17 were induced from primary peripheral blood CD4+ T cells upon stimulation and they can induce stromal cells to produce proinflammatory and hematopoietic cytokines.