

Recombinant Human Thymic Stromal Lymphopoietin (rHuTSLP)

PrimeGene Technical Data Sheet

Catalog Number:	102-07
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 15.1 kDa, a single non-glycosylated polypeptide chain containing 132 amino acids.
Quantity:	2µg/10µg/1000µg
AA Sequence:	MYDFTNCDFE KIKAAAYLSTI SKDLITYMSG TKSTEFNNTV SCSNRPHCLT EIQLSLTFNPT AGCASLAKEM FAMKTKAALA IWCPGYSETQ INATQAMKKR RKRKVTTNKC LEQVSQLQGL WRRFNRPLLK QQ
Purity:	> 98 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell proliferation assay using human IL-7Rα and human TSLP R co-transfected murine BaF3 pro-B cells is less than 0.3 ng/ml, corresponding to a specific activity of > 3.3 × 10 ⁶ IU/mg.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM NaCl.
Endotoxin:	Less than 1 EU/µg of rHuTSLP as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Human Thymic Stromal Lymphopoietin

Thymic stromal lymphopoietin (TSLP) is a hemopoietic protein belonging to the cytokine family and is known to play an important role in the maturation of T cell populations through activation of antigen presenting cells. It is mainly expressed in a number of tissues including heart, liver and prostate. TSLP signals through a heterodimeric receptor complex composed of the thymic stromal lymphopoietin receptor and the IL-7R alpha chain. After binding STAT5 phosphorylation is induced resulting in the expression of downstream transcription factors. Like IL-7, TSLP induces phosphorylation of STAT3 and STAT5, but uses kinases other than the JAKs for activation. TSLP has the functions of enhancing the maturation of CD11c+ dendritic cells and inducing allergic inflammation by directly activating mast cells. Its expression is linked to many disease states including asthma, inflammatory arthritis, atopic dermatitis, and eczema and other allergic states. But the factors inducing the activation of TSLP release are not clearly defined. Human TSLP shares approximately 43 % a.a. sequence identity with mouse TSLP.