

**Recombinant Human Tumor Necrosis  
Factor-alpha/TNFSF2, His  
(rHuTNF- $\alpha$ /TNFSF2, His)  
PrimeGene Technical Data Sheet**

---

<b>Catalog Number:</b>	103-01H
<b>Source:</b>	<i>Escherichia coli</i> .
<b>Molecular Weight:</b>	Approximately 18.3 kDa, a single, non-glycosylated polypeptide chain containing 164 amino acids with Met and 6 $\times$ His at N-terminus.
<b>Quantity:</b>	10 $\mu$ g/50 $\mu$ g/1000 $\mu$ g
<b>AA Sequence:</b>	<b>MHHHHHHVRS SSRTPSDKPV AHVVANPQAE GQLQWLNRRR NALLANGVEL RDNQLVVPSE GLYLIYSQVL FKGQGCPSTH VLLTHTISRI AVSYQTKVNL LSAIKSPCQR ETPEGAEAKP WYEPIYLGGV FQLEKGDRLS AEINRPDYLD FAESGQVYFG IIAL</b>
<b>Purity:</b>	> 97 % by SDS-PAGE and HPLC analyses.
<b>Biological Activity:</b>	Fully biologically active when compared to standard. The ED <sub>50</sub> as determined by a cytotoxicity assay using murine L929 cells is less than 0.05 ng/ml, corresponding to a specific activity of > 2.0 $\times$ 10 <sup>7</sup> IU/mg in the presence of actinomycin D.
<b>Physical Appearance:</b>	Sterile Filtered White lyophilized (freeze-dried) powder.
<b>Formulation:</b>	Lyophilized from a 0.2 $\mu$ m filtered concentrated solution in PBS, pH 7.0.
<b>Endotoxin:</b>	Less than 1 EU/ $\mu$ g of rHuTNF- $\alpha$ /TNFSF2, His as determined by LAL method.
<b>Reconstitution:</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at $\leq$ -20 °C. Further dilutions should be made in appropriate buffered solutions.
<b>Shipping:</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage:</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"><li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li><li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li><li>● 3 months, -20 to -70 °C under sterile conditions after reconstitution.</li></ul>
<b>Usage:</b>	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. <b>NOT FOR HUMAN USE.</b>

---

### ***Human Tumor Necrosis Factor-alpha/TNFSF2***

Tumor necrosis factor alpha (TNF- $\alpha$ ) is the best-know member of the TNF-family, which can cause cell death. This protein is produced by neutrophils, activated lymphocytes, macrophages, NK cells, LAK cells, astrocytes endothelial cells, smooth muscle cells and some transformed cells. TNF- $\alpha$  occurs as a secreted, soluble form and as a membrane-anchored form, both of which are biologically active. The naturally-occurring form of TNF- $\alpha$  is glycosylated, but non-glycosylated recombinant TNF- $\alpha$  has comparable biological activity. The biologically active native form of TNF- $\alpha$  is reportedly a trimer. Human and murine TNF- $\alpha$  show approximately 79 % homology at the amino acid level and cross-reactivity between the two species. Two types of receptors for TNF- $\alpha$  have been described and virtually all cell types studied show the presence of one or both of these receptor types.