

Recombinant Human LR3 Insulin-like Growth Factor-1 (rHuLR3 IGF-1)

PrimeGene Technical Data Sheet

Catalog Number:	105-03
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 9.1 kDa, a single non-glycosylated polypeptide chain containing 83 amino acids.
Quantity:	1mg/5mg/20mg/50mg
AA Sequence:	MFPAMPLSSL FVNGPRTLPG AELVDALQFV CGDRGFYFNK PTGYGSSRR APQTGIVDEC CFRSCDLRRL EMYCAPLKPA KSA
Purity:	>98 % by SDS-PAGE analyses. >90 % by RP-HPLC analyses.
Biological Activity:	Assay 1: Fully biologically active when compared to standard. Measured in a serum-free cell proliferation assay using human MCF-7 cells. The ED ₅₀ for this effect is 0.3-1.5 ng/ml, corresponding to a specific activity of > 6.7 × 10 ⁵ U/mg. Assay 2: Fully biologically active when compared to standard. The ED ₅₀ as determined by the stimulation of protein synthesis using rat L6 myoblasts is less than 10 ng/ml, corresponding to a specific activity of > 1.0 × 10 ⁵ U/mg.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 μm filtered concentrated solution in 20 mM PB, pH 7.2.
Endotoxin:	Less than 0.01 EU/μg of rHuLR3 IGF-1 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● Refer to lot specific COA for the Use by Date when stored at 2 to 8 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Human LR3 Insulin-like Growth Factor-1

IGF-1 belonged to the insulin gene family, is a mitogenic polypeptide growth factor that stimulate the proliferation and survival of various cell types including muscle, bone, and cartilage tissue in vitro. It is produced primarily by the liver as an endocrine hormone as well as in target tissues in a paracrine/autocrine fashion. The production of IGF-1 is stimulated by growth hormone (GH) and can be retarded by undernutrition, growth hormone insensitivity, lack of growth hormone receptors, or failures of the downstream signaling pathway post GH receptor including SHP2 and STAT5B. The LR3 IGF-1 is a long-term analog of human IGF-1, specifically designed and manufactured for mammalian cell culture to support large-scale manufacturing of recombinant biopharmaceuticals.