

Recombinant Human Interferon-alpha1b (rHuIFN-a1b)

PrimeGene Technical Data Sheet

Catalog Number: 106-02

Source: Escherichia coli.

Molecular Weight: Approximately 19.5 kDa, a single non-glycosylated polypeptide chain containing 167 amino acids.

Quantity: $2\mu g/10\mu g/1000\mu g$

AA Sequence: MCDLPETHSL DNRRTLMLLA QMSRISPSSC LMDRHDFGFP QEEFDGNQFQ

> KAPAISVLHE LIQQIFNLFT TKDSSAAWDE DLLDKFCTEL YQQLNDLEAC VMQEERVGET PLMNVDSILA VKKYFRRITL YLTEKKYSPC AWEVVRAEIM

RSLSLSTNLQ ERLRRKE

Purity: > 96 % by SDS-PAGE and HPLC analyses.

Biological Activity: Fully biologically active when compared to standard. The specific activity determined by an anti-viral

assay is no less than 1.0×10^8 IU/mg.

Sterile Filtered White lyophilized (freeze-dried) powder. Physical Appearance:

Formulation: Lyophilized from a 0.2 µm filtered solution in PBS, pH 7.4, containing 4 % mannitol and 1 % HSA.

Endotoxin: Less than 1 EU/μg of rHuIFN-α1b as determined by LAL method.

Reconstitution: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the

> bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and

stored at \leq -20 °C. Further dilutions should be made in appropriate buffered solutions.

Shipping: The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature

recommended below.

Stability & Storage: Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

12 months from date of receipt, -20 to -70 °C as supplied.

1 month, 2 to 8 °C under sterile conditions after reconstitution.

3 months, -20 to -70 °C under sterile conditions after reconstitution.

Usage: This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further

evaluation purposes. NOT FOR HUMAN USE.

Human Interferon-alpha1b

IFN-αs are proteins secreted by leukocyte. They are mainly involved in innate immune response against viral infection. The IFNα family has 13 subtypes and 23 different variants. The individual proteins have molecular masses between 19-26 kDa and consist of proteins with lengths of 156-166 and 172 amino acids. All IFN-α subtypes possess a common conserved sequence region between amino acid positions 115-151 while the amino-terminal ends are variable. Many IFN-alpha subtypes differ in their sequences at only one or two positions. Naturally occurring variants also include proteins truncated by 10 amino acids at the carboxy-terminal end.

Rev. 08/20/2018 V.3

Email: info.pg@bio-techne.com

Shanghai PrimeGene Bio-Tech Co., Ltd.

Website: www.primegene.com.cn Tel: +86 21 52380373

Website: www.primegene.com

Fax: +86 21 61077348