

## PrimeGene Technical Data Sheet

---

<b>Catalog Number:</b>	144-09
<b>Source:</b>	<i>Escherichia coli</i> .
<b>Molecular Weight:</b>	Approximately 23.1 kDa, a single non-glycosylated polypeptide chain containing 205 amino acids.
<b>Quantity:</b>	5µg/20µg/1000µg
<b>AA Sequence:</b>	LGEVGSYFGV QDAVPFGNVP VLPVDSPVLL SDHLGQSEAG GLPRGPAVTD LDHLKGILRR RQLYCRTGFH LEIFPNGTIQ GTRKDHSRFG ILEFISIAVG LVSIRGVDSG LYLGMNEKGE LYGSEKLTQE CVFREQFEEN WYNTYSSNLY KHVDTGRRYY VALNKDGT EGTRTKRHQK FTHFLRPVD PDKVPELYKD ILSQS
<b>Purity:</b>	> 95 % by SDS-PAGE and HPLC analyses.
<b>Biological Activity:</b>	Fully biologically active when compared to standard. The ED <sub>50</sub> as determined by thymidine uptake assay using FGF-receptors transfected BaF3 cells is less than 0.5 ng/ml, corresponding to a specific activity of > 2.0 × 10 <sup>6</sup> IU/mg.
<b>Physical Appearance:</b>	Sterile Filtered White lyophilized (freeze-dried) powder.
<b>Formulation:</b>	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM Tris, 400 mM NaCl, pH 8.0.
<b>Endotoxin:</b>	Less than 1 EU/µg of rRtFGF-9-1 as determined by LAL method.
<b>Reconstitution:</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
<b>Shipping:</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage:</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"><li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li><li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li><li>● 3 months, -20 to -70 °C under sterile conditions after reconstitution.</li></ul>
<b>Usage:</b>	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. <b>NOT FOR HUMAN USE.</b>

---

### ***Rat Fibroblast Growth Factor-9***

Fibroblast growth factor-9 (FGF-9) is a member of the fibroblast growth factor (FGF) family. All FGF family members are heparin binding growth factors with a core 120 amino acid (a.a.) FGF domain that allows for a common tertiary structure. FGF-9 plays an important role in the regulation of embryonic development, cell proliferation, cell differentiation and cell migration. This protein was isolated as a secreted factor that exhibits a growth-stimulating effect on cultured glial cells. FGF-9 is a monomer and interacts with FGFR1, FGFR2, FGFR3 and FGFR4. The rat FGF-9 shares 98 % a.a. sequence identity with human FGF-9.