

# Recombinant Rat Resistin (rRtResistin)

## PrimeGene Technical Data Sheet

---

<b>Catalog Number:</b>	14A-03
<b>Source:</b>	<i>Escherichia coli</i> .
<b>Molecular Weight:</b>	Approximately 20.2 kDa, a disulfide-linked homodimer consisting of two 95 amino acid polypeptide chains.
<b>Quantity:</b>	5µg/25µg/1000µg
<b>AA Sequence:</b>	MPSMSLCPMD EAISKKINQD FSSLLPAAMK NTVLHCWSVS SRGRLASCPE GTTVTSCSCG SGCGSWDVRE DTMCHCQCGS IDWTAARCCT LRVGS
<b>Purity:</b>	> 97 % by SDS-PAGE and HPLC analyses.
<b>Biological Activity:</b>	Test in process.
<b>Physical Appearance:</b>	Sterile Filtered White lyophilized (freeze-dried) powder.
<b>Formulation:</b>	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH7.4, with 0.02 % Tween-20.
<b>Endotoxin:</b>	Less than 0.1 EU/µg of rRtResistin as determined by LAL method.
<b>Reconstitution:</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
<b>Shipping:</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage:</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"><li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li><li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li><li>● 3 months, -20 to -70 °C under sterile conditions after reconstitution.</li></ul>
<b>Usage:</b>	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. <b>NOT FOR HUMAN USE.</b>

---

### ***Rat Resistin***

Resistin, also known as adipocyte-specific secretory factor (ADSF) and found in inflammatory zone 3 (FIZZ3), is an adipocyte-derived protein. Mature rat and human Resistin share 54% amino acid sequence identity. Northern blot analysis shows that Resistin is expressed in adipose tissue, and its expression is markedly induced during adipocyte differentiation of 3T3-L1 preadipocyte cells.