

PrimeGene Technical Data Sheet

Catalog Number:	203-01
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 8.6 kDa, a single non-glycosylated polypeptide chain containing 76 amino acids and comprises only the chemokine domain of Human Fractalkine.
Quantity:	5µg/20µg/100µg
AA Sequence:	QHHGVTKCNI TCSKMTSKIP VALLIHYQQN QASCGKRAII LETRQHRLFC ADPKEQWVKD AMQHLDLRQAA ALTRNG
Purity:	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human T-lymphocytes is in a concentration of 5.0-10 ng/ml.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 50 mM NaCl.
Endotoxin:	Less than 1 EU/µg of rHuFractalkine/CX3CL1 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Human Fractalkine/CX3CL1

CX3CL1 recently identified through bioinformatics is the only known member of the CX3C chemokine family and it is also commonly known under the names fractalkine (in humans) and neurotactin (in mice). Unlike other known chemokines, CX3CL1 is a type 1 membrane protein containing a chemokine domain tethered on a long mucinlike stalk. The soluble form of CX3CL1 is chemotactic for T-cells and monocytes, but not for neutrophils. In addition, it may play a role in regulating leukocyte adhesion and migration processes at the endothelium. Recombinant Human CX3CL1 which is a single non-glycosylated polypeptide chains contains 76 amino acids and it shares approximately 78 % and 83 % amino acid sequence homology with the murine and rat protein.