

PrimeGene Technical Data Sheet

Catalog Number:	204-01
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 8.6 kDa, a single, non-glycosylated polypeptide chain containing 74 amino acids.
Quantity:	2µg/10µg/1000µg
AA Sequence:	SKSMQVPFSR CCFSFAEQEI PLRAILCYRN TSSICSNEGL IFKLRGKEA CALDTVGWVQ RHRKMLRHCP SKRK
Purity:	>97% by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human T-lymphocytes is in a concentration range of 10-100 ng/ml.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 100 mM NaCl.
Endotoxin:	Less than 1 EU/µg of rHuI-309/CCL1 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Human I-309/CCL1

Chemokine (C-C motif) ligand 1 (CCL1) belongs to a family inflammatory cytokines and also known as chemokines. It is a small glycoprotein secreted by activated T cells with a molecular weight of approximately 8.5 kDa. CCL1 attracts monocytes, NK cells, and immature B cells and dendritic cells by interacting with a cell surface chemokine receptor called CCR8. Human CCL1 has been assumed to be a homologue of the mouse TCA3. While the two proteins share only approximately 42 % amino acid sequence identity, both chemokines contain an extra pair of cysteine residues not found in most other chemokines.