

**Recombinant Human Monocyte Chemotactic  
Protein-4/CCL13  
(rHuMCP-4/CCL13)  
PrimeGene Technical Data Sheet**

---

<b>Catalog Number:</b>	204-13
<b>Source:</b>	<i>Escherichia coli</i> .
<b>Molecular Weight:</b>	Approximately 8.6 kDa, a single non-glycosylated polypeptide chain containing 75 amino acids.
<b>Quantity:</b>	5µg/20µg/1000µg
<b>AA Sequence:</b>	QPDALNVPST CCFTFSSKKI SLQRLKSYVI TTSRCPQKAV IFRTKLGKEI CADPKEKWVQ NYMKHLGRKA HTLKT
<b>Purity:</b>	> 96 % by SDS-PAGE and HPLC analyses.
<b>Biological Activity:</b>	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human monocytes is in a concentration of 10-100 ng/ml.
<b>Physical Appearance:</b>	Sterile Filtered White lyophilized (freeze-dried) powder.
<b>Formulation:</b>	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 130 mM NaCl.
<b>Endotoxin:</b>	Less than 1 EU/µg of rHuMCP-4/CCL13 as determined by LAL method.
<b>Reconstitution:</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20°C. Further dilutions should be made in appropriate buffered solutions.
<b>Shipping:</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage:</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"><li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li><li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li><li>● 3 months, -20 to -70 °C under sterile conditions after reconstitution.</li></ul>
<b>Usage:</b>	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. <b>NOT FOR HUMAN USE.</b>

---

***Human Monocyte Chemotactic Protein-4/CCL13***

Human CCL13 is belonging to the CC chemokine family and is encoded by the gene CCL13 in humans. CCL13 (MCP-4) shares 56-61 % sequence identity with MCP-1 (CCL2) and MCP-3 (CCL7) and is 60 % identical to Eotaxin (CCL11). CCL13 was a potent chemoattractant for monocytes and eosinophils and stimulated histamine release from basophils. CCL13 can induce a calcium flux in HEK-293 cells transfected with the receptor CCR2B and CCR3. That shows the function receptors of CCL3 are CCR2B and CCR3.