

**Recombinant Rat Neutrophil Activating
Peptide 2/CXCL7
(rRtNAP-2/CXCL7)
PrimeGene Technical Data Sheet**

Catalog Number:	241-07
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 6.8 kDa, a single non-glycosylated polypeptide chain containing 62 amino acids.
Quantity:	2µg/10µg/1000µg
AA Sequence:	IELRCRCTNT LSGIPLNSIS RVNVFRPGAH CDNVEVIATL KNGKEVCLDP TAPMIKKIVK KI
Purity:	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The biologically active determined by a chemotaxis bioassay using human CXCR2 transfected murine BaF3 cells is in a concentration range of 0.1-1.0 ng/ml.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Endotoxin:	Less than 1 EU/µg of rRtNAP-2/CXCL7 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Rat Neutrophil Activating Peptide 2/CXCL7

CXCL7 is a small cytokine which is also an isoform of Beta-Thromboglobulin or Pro-Platelet basic protein belonging to the CXC chemokine family and it is released in large amounts from platelets following their activation. CXCL7 stimulates DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E2 secretion, and synthesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by human synovial cells. Recombinant Rat CXCL7 contains 62 amino acids which is a single non-glycosylated polypeptide chain. In addition, The rat CXCL7 shares 58 % and 77 % a.a. sequence identity with human and mouse CXCL7.